



# Nutrient salvaging and metabolism by the intracellular pathogen *Legionella pneumophila*

Maris V. Fonseca<sup>1</sup> and Michele S. Swanson<sup>2\*</sup>

<sup>1</sup> Science and Mathematics Division, Monroe County Community College, Monroe, MI, USA

<sup>2</sup> Department of Microbiology and Immunology, University of Michigan Medical School, Ann Arbor, MI, USA

## Edited by:

Monique Barel, Institut National de la Santé et de la Recherche Médicale, France

## Reviewed by:

Monique Barel, Institut National de la Santé et de la Recherche Médicale, France  
Max Maurin, Assistance Publique-Hôpitaux de Marseille, Université Aix-Marseille II, France

## \*Correspondence:

Michele S. Swanson, Department of Microbiology and Immunology, University of Michigan Medical School, 6733 Medical Science Building II, 1150 West Medical Center Drive, Ann Arbor, MI 48109-5620, USA  
e-mail: mswanson@umich.edu

The Gram-negative bacterium *Legionella pneumophila* is ubiquitous in freshwater environments as a free-swimming organism, resident of biofilms, or parasite of protozoa. If the bacterium is aerosolized and inhaled by a susceptible human host, it can infect alveolar macrophages and cause a severe pneumonia known as Legionnaires' disease. A sophisticated cell differentiation program equips *L. pneumophila* to persist in both extracellular and intracellular niches. During its life cycle, *L. pneumophila* alternates between at least two distinct forms: a transmissive form equipped to infect host cells and evade lysosomal degradation, and a replicative form that multiplies within a phagosomal compartment that it has retooled to its advantage. The efficient changeover between transmissive and replicative states is fundamental to *L. pneumophila*'s fitness as an intracellular pathogen. The transmission and replication programs of *L. pneumophila* are governed by a number of metabolic cues that signal whether conditions are favorable for replication or instead trigger escape from a spent host. Several lines of experimental evidence gathered over the past decade establish strong links between metabolism, cellular differentiation, and virulence of *L. pneumophila*. Herein, we focus on current knowledge of the metabolic components employed by intracellular *L. pneumophila* for cell differentiation, nutrient salvaging and utilization of host factors. Specifically, we highlight the metabolic cues that are coupled to bacterial differentiation, nutrient acquisition systems, and the strategies utilized by *L. pneumophila* to exploit host metabolites for intracellular replication.

**Keywords:** *Legionella pneumophila*, differentiation, intracellular metabolism, metabolic cues, phagosomal transporters


## INTRODUCTION

The facultative intracellular bacterium *Legionella pneumophila* parasitizes protozoa in aquatic environments and alveolar macrophages in susceptible human hosts. *L. pneumophila* survives in nature by virtue of a differentiation cycle in which distinct cell types interconvert in response to environmental and metabolic fluctuations. In its planktonic transmissive form, *L. pneumophila* is motile, resistant to multiple environmental stresses, including nutrient starvation, and infectious to host cells (Rowbotham, 1983, 1986; Byrne and Swanson, 1998). In the transmissive phase, effectors translocated across the bacterial membrane and virulence factors on the *L. pneumophila* surface arrest phagosome maturation to establish a replication vacuole derived from the host's endoplasmic reticulum (Swanson and Isberg, 1995; Byrne and Swanson, 1998; Joshi et al., 2001; Fernandez-Moreira et al., 2006; Ensminger and Isberg, 2009; Isberg et al., 2009; Rolando and Buchrieser, 2012; Amyot et al., 2013; **Figure 1**). The replicative form of *L. pneumophila* multiplies intracellularly within such vacuoles, which in some host cells mature into acidic lysosomal vacuoles that support bacterial growth (Sturgill-Koszycki and Swanson, 2000; Xu et al., 2010).

*L. pneumophila* can also differentiate to other cell types that remain to be characterized in molecular detail. During its life cycle inside protozoa, HeLa cells, epithelial cells or even

clinical specimens, transmissive *L. pneumophila* differentiate further to a mature intracellular form (MIF), a highly infectious, metabolically-resting, cyst-like form that is observed late during infection (Faulkner and Garduno, 2002; Garduno et al., 2002; Greub and Raoult, 2003; Faulkner et al., 2008; Garduno, 2008). The resilience of extracellular *L. pneumophila* is further evidenced by its survival in a viable but non-culturable (VBNC) state that can arise when either transmissive or stationary phase *L. pneumophila* or MIFs are exposed to severe conditions in water environments (Steinert et al., 1997; Garduno, 2008; Al-Bana et al., 2014). *L. pneumophila* can also form monospecies biofilms or colonize multi-species biofilm communities (Abdel-Nour et al., 2013).

The ability to exploit extracellular and intracellular niches and endure environmental stresses, including nutritional starvation, equips *L. pneumophila* to persist in nature. Indeed, adaptation to the highly variable conditions encountered by *L. pneumophila* requires swift morphogenetic and physiological transformations (Byrne and Swanson, 1998; Garduno et al., 2002; Molofsky and Swanson, 2004; Garduno, 2008). Accordingly, the *L. pneumophila* life cycle is controlled by multipronged regulatory systems that control gene expression; these include a variety of transcriptional regulatory proteins, two-component systems, non-coding RNA (ncRNA) molecules, the stringent



response pathway, and metabolites (Hammer and Swanson, 1999; Bachman and Swanson, 2001, 2004; Hammer et al., 2002; Molofsky and Swanson, 2003; Molofsky et al., 2005; Hovel-Miner et al., 2009; Dalebroux et al., 2009, 2010; Edwards et al., 2009, 2010; Sahr et al., 2009; Albert-Weissenberger et al., 2010).

Whether in extracellular or intracellular environments, differentiation of transmissible *L. pneumophila* to the replicative form is tightly coupled to its metabolic state (Hammer and Swanson, 1999; Sauer et al., 2005; Dalebroux et al., 2009; Edwards et al., 2009, 2010). When transmissible *L. pneumophila* encounter abundant nutrients in their environment, such as amino acids, these metabolites trigger differentiation to the replicative form (Hammer and Swanson, 1999; Sauer et al., 2005; Figure 1). As *L. pneumophila* multiplies, nutrient consumption or accumulation of certain metabolites alters the bacterium's nutritional milieu. The metabolic changes that these alterations provoke are relayed to regulatory components that include the stringent response, the LetA/S two-component system and alternative sigma factors to trigger differentiation of *L. pneumophila* into the transmissible form (Edwards et al., 2009, 2010; Figure 1). The transmissible form of *L. pneumophila* is suited to seek a new niche that supports replication. Thus, metabolic cues serve not only as gauges of fitness of the environment for replication but also as triggers of differentiation into other cell types, such as the transmissible form.

The sensing of metabolic signals to adopt a particular cell type does not constitute the only strategy intracellular *L. pneumophila* utilizes to meet its physiological and metabolic needs. Among the many effector molecules translocated by the Icm/Dot type IVB secretion system are proteins that exploit host pathways either by altering eukaryotic phospholipid metabolism or by utilizing the host proteasome to generate an amino acid supply (Al-Quadan et al., 2012; Viner et al., 2012; Bruckert et al., 2014). As another means to exploit the host environment for replication,

intracellular *L. pneumophila* employs components of its own metabolism such as nutrient salvaging systems (Cianciotto et al., 2005; Sauer et al., 2005; Yip et al., 2011; Fonseca et al., 2014). In this review, we showcase the growing body of knowledge of the metabolic components derived from both bacterial and host sources that contribute to *L. pneumophila* replication within host cells.

## IN VITRO EXPERIMENTAL PROBES OF INTRACELLULAR METABOLISM BY *Legionella pneumophila*

The study of *L. pneumophila* metabolism, particularly of intracellular forms, has been expedited by a number of genetic and molecular tools and *in vitro* studies of metabolism. Of particular importance has been the ability to generate mutants using various methodologies, including enrichments (Berger and Isberg, 1993; Swanson and Isberg, 1996), transposon libraries (Hammer and Swanson, 1999; Sauer et al., 2005), natural competence (Sexton and Vogel, 2004; Buchrieser and Charpentier, 2013), and recombineering (Bryan et al., 2011). Using these techniques, *L. pneumophila* mutants defective at particular stages of the intracellular life cycle have been isolated and characterized to gain insight into both the fate of *L. pneumophila* in phagocytes and the host components that support infection.

The ability to study the growth-phase dependent expression of panels of replicative and transmissible traits of *L. pneumophila* in synchronous broth cultures has also greatly advanced our understanding of the regulation of this bacterium's life cycle and its metabolism (Rowbotham, 1986; Byrne and Swanson, 1998; Molofsky and Swanson, 2004; Dalebroux et al., 2009; Edwards et al., 2009). This knowledge has enhanced our ability to assess the requirement of specific functions for particular cell types *in vivo* (Sauer et al., 2005; Dalebroux et al., 2010). For instance, *L. pneumophila* mutants of the phagosomal transporter A (*phtA*) display an intracellular growth defect that can be bypassed *in vivo* by threonine supplementation (Sauer et al., 2005). Moreover, without threonine supplementation, *phtA* mutants are locked in the transmissible form and fail to differentiate into the replicative form inside macrophages. Similarly, the reciprocal expression of replicative and transmissible traits can be utilized to assess the requirement of various gene products for bacterial differentiation in the host (Hammer et al., 2002; Molofsky and Swanson, 2003; Dalebroux et al., 2010).

The response of *L. pneumophila* to particular metabolic cues has been probed by coupling various molecular methods to assays of growth-phase dependent traits (Hammer and Swanson, 1999; Edwards et al., 2009). These include the use of transcriptional fusions (Hammer and Swanson, 1999; Hammer et al., 2002; Edwards et al., 2009), phenotype microarrays (Sauer et al., 2005; Edwards et al., 2009) and transcriptional profile studies (Bruggemann et al., 2006; Dalebroux et al., 2010; Edwards et al., 2010; Hovel-Miner et al., 2010; Faucher et al., 2011). These tools have facilitated experiments to probe how intracellular *L. pneumophila* sense its nutritional milieu in macrophages and adopt the appropriate cell fate (Dalebroux et al., 2009, 2010).

Insights into the nutritional requirements of *L. pneumophila* in host cells have also been obtained from classical nutritional studies using broth cultures. Groundbreaking experiments revealed

key nutritional requirements of *L. pneumophila*: the absolute requirement for oxygen for its metabolism, the use of amino acids as major sources of carbon and energy (Pine et al., 1979; Reeves et al., 1981; Ristroph et al., 1981; Tesh et al., 1983), the requirements for specific nutrients in host cells (Sauer et al., 2005), the utilization of host-derived nutrients (Al-Quadani et al., 2012; Bruckert et al., 2014), and the identification of nutritional auxotrophies (Pine et al., 1979; George et al., 1980; Ristroph et al., 1981; Tesh and Miller, 1981). In other instances, the ability to assess the nutritional requirements of *L. pneumophila* in broth has allowed genes of unknown function to be linked to particular metabolic processes (Sauer et al., 2005; Fonseca et al., 2014).

Heterologous genetic systems have enabled the function of several *L. pneumophila* proteins to be deduced. This approach proved instrumental in gathering supporting evidence for *in silico* analyses and observed intracellular growth phenotypes (Molofsky and Swanson, 2003; Bachman and Swanson, 2004; Nasrallah et al., 2011; Viner et al., 2012; Fonseca et al., 2014).

Isotopologue profiling has been implemented as a tool to probe the metabolic pathways of assimilation of various precursors into building blocks in *L. pneumophila* (Eylert et al., 2010). First, the *in vitro* isotopologue studies confirmed several of the amino acid auxotrophies that had been observed in broth cultures of *L. pneumophila*, namely those of methionine, arginine, valine, leucine, and isoleucine. Furthermore, by using isotopologues of glucose and the amino acid serine, the investigators determined that *L. pneumophila* can metabolize glucose using the Entner-Doudoroff (ED) pathway and that carbon from glucose is primarily used for the synthesis of poly-3-hydroxybutyrate, a storage compound that *L. pneumophila* utilizes under nutrient starvation conditions (James et al., 1999; Garduno et al., 2002). The contribution of the ED pathway was tested by generation of a *L. pneumophila* strain Paris mutant in the putative glucose-6-phosphate-1-dehydrogenase gene (*zwf*) (Eylert et al., 2010). In multiple-round co-infection experiments, the *zwf* mutant was outcompeted by wild type after >2 weeks, pointing to a contribution of the *zwf* gene in fitness during prolonged co-culture with amoebae. How the *zwf* gene in particular and the ED pathway in general contribute to *Legionella* fitness in the environment requires further investigation. Thus, implementation of isotopologue labeling techniques has increased our understanding of *L. pneumophila* metabolism and has led to further development of this methodology to study the intracellular metabolism of *L. pneumophila* (Heuner and Eisenreich, 2013).

Our current knowledge of *L. pneumophila* metabolism relies on the integration of various *in vitro* and *in vivo* approaches. A caveat of incorporating the results of various experimental approaches to understanding *L. pneumophila* physiology is that the metabolic state of the bacterium may vary widely depending on the cell type, growth culture conditions, and sensitivity of the assays employed, which may lead to apparent discrepancies in experimental results.

## REGULATION OF THE *L. pneumophila* LIFE CYCLE BY METABOLITES

A salient feature of bacterial cells is their ability to adapt to constant physicochemical changes in their environment. The tuning

of bacterial physiology by controlling gene expression at the transcriptional and translational level has been the object of intense study (Henkin, 2000; Browning and Busby, 2004; Serganov and Patel, 2012). In addition to gene regulation, bacteria ensure energy efficiency and persistence in nature by employing another strategy, one perhaps less appreciated by molecular biologists: metabolic control. The latter mainly consists of post-translational control of metabolism by metabolites. Mechanisms of post-translational control common in biology include allosteric control and feedback inhibition (Sanwall, 1970; LaRossa et al., 1987; Allen et al., 2002; Starai and Escalante-Semerena, 2004; Wolfe, 2005; Bazaruto and Downs, 2014). The overlap between many of these genetic and metabolic regulatory machineries further supports how bacterial physiology is subject to sophisticated control. The collective contributions of these studies have been invaluable in our current understanding of the factors that allow bacterial adaptation to a wide variety of alterations in their surroundings.

That the *L. pneumophila* intracellular life cycle is also subject to regulation by metabolites is a fact that is supported by numerous lines of evidence (Byrne and Swanson, 1998; Hammer and Swanson, 1999; Sauer et al., 2005; Dalebroux et al., 2009; Edwards et al., 2009; Hovel-Miner et al., 2010). In several examples we shall discuss here, it is clear that a subset of metabolites exert their effect, at least in part, at the transcriptional level. Whether various metabolites govern the *L. pneumophila* life cycle by mechanisms such as allostery is an intriguing question that requires further experimentation, particularly, biochemical analyses.

## AMINO ACIDS

The preference of *L. pneumophila* for amino acids as main sources of carbon and energy was first deduced from experiments in broth cultures (Pine et al., 1979; Reeves et al., 1981; Ristroph et al., 1981; Tesh et al., 1983). More recently, isotopologue studies further indicated that the amino acid serine supports carbon metabolism in *L. pneumophila* as well as confirmed some of the amino acid auxotrophies that had been determined previously using broth cultures (Pine et al., 1979; George et al., 1980; Ristroph et al., 1981; Tesh and Miller, 1981; Eylert et al., 2010). These and other amino acid auxotrophies had also been suggested by *in silico* genomic analyses (Fonseca et al., 2008). Moreover, as exemplified by the characterization of *phtA* mutants in macrophages, differentiation of transmissive *L. pneumophila* to the replicative form responds to cues that indicate amino acid abundance. Thus, a plentiful amino acid supply promotes replication of either intracellular or extracellular *L. pneumophila*.

Fundamental to our current understanding of the *L. pneumophila* life cycle has been the observation that, when starved for amino acids, *L. pneumophila* switches from the replicative form to the transmissive form. Transmissive *L. pneumophila* express a number of virulence traits such as motility, lysosome evasion and cytotoxicity (Byrne and Swanson, 1998; Hammer and Swanson, 1999; Bachman and Swanson, 2001; Lynch et al., 2003) that promote escape from a spent host and infection of a naïve one. The response to amino acid starvation is coordinated by synthesis of the alarmone ppGpp by the RelA synthetase (Hammer and Swanson, 1999; Dalebroux et al., 2009, 2010). The ppGpp signal



transducer in turn induces expression of a series of genes typical of stress responses in bacteria (Gentry et al., 1993; Durfee et al., 2008; Traxler et al., 2008; Zhao and Houry, 2010). Indeed, ppGpp triggers activation of various regulatory components in *L. pneumophila*, such as alternative sigma factors (Rasis and Segal, 2009; Sahr et al., 2009; Edwards et al., 2010), in a complex regulatory cascade that also requires the LetA/LetS two-component system (Hammer et al., 2002; Dalebroux et al., 2010). The dramatic morphogenetic and physiological alterations exhibited by *L. pneumophila* in response to amino acid scarcity or abundance underscore the importance of these nutrients to its life cycle and virulence.

The ability of amino acids to interact with regulatory components that are pivotal for intracellular replication by *L. pneumophila* was further evidenced in a study of arginine metabolism in this bacterium (Hovel-Miner et al., 2010). The study stemmed from the experimental observation that mutations in the arginine repressor (ArgR) protein of *L. pneumophila* confer an intracellular replication defect within host *Acanthamoeba castellanii* (Hovel-Miner et al., 2009). ArgR protein represses genes required for *de novo* arginine biosynthesis from glutamate (Maas, 1994). Whereas *L. pneumophila* lacks many of the genes that other species require for early steps in arginine biosynthesis, the bacterium can synthesize arginine in defined medium supplemented with either of the precursors ornithine or citrulline (Tesh et al., 1983; Hovel-Miner et al., 2010). Hovel-Miner and colleagues demonstrated that *L. pneumophila* lacking the genes for either a putative amino acid ABC transporter or the last step of the arginine biosynthetic route fail to grow in defined medium unless provided with arginine. Nevertheless, both mutant strains do grow intracellularly, and their growth was not stimulated by addition of arginine to the infection buffer. Accordingly, it was deduced that *L. pneumophila* can tolerate arginine auxotrophy because the pathogen obtains arginine from the host to support intracellular replication. The latter finding was intriguing considered in the context of the intracellular growth defect conferred by *argR* mutants in amoeba. Global transcriptional analyses employed to address this paradox revealed that in *L. pneumophila* the ArgR repressor regulates a number of components apparently unrelated to arginine metabolism but required for intracellular replication.

Among the components regulated by ArgR in *L. pneumophila* are genes encoding proteins of the Icm/Dot Type IV secretion system (subject to negative regulation) and its translocated substrates (subject to positive regulation). These data led to the proposal that arginine and ArgR are required for assembly of the Icm/Dot system at the appropriate time during the *L. pneumophila* life cycle and for the synthesis of functions that promote establishment of the *L. pneumophila* replication vacuole.

Noteworthy is the fact that in organisms like *Escherichia coli* ArgR exerts its regulatory effect with arginine as a co-repressor (Lim et al., 1987; Maas, 1994). The results of the global transcriptional analyses suggest that arginine has a similar role in *L. pneumophila* (Hovel-Miner et al., 2010). This finding together with the similarity of amino acid sequences and DNA-binding sites of the ArgR protein in *L. pneumophila* to ArgR proteins of *E. coli* and *Bacillus subtilis* further highlight the importance of

amino acid metabolic cues to the regulatory circuits that control intracellular replication and differentiation in *L. pneumophila*.

## FATTY ACIDS

Short-chain fatty acids (SCFAs) are known regulators of metabolism and virulence in bacteria. For example, in *Salmonella enterica* serovar Typhimurium, exposure to SCFAs like acetate and propionate can have deleterious effects on cell metabolism. Among these are effects on acyl-CoA homeostasis, depletion of free CoA pools, and accumulation of toxic metabolic intermediates, which have also been observed in other bacteria like *E. coli* and *Rhodobacter sphaeroides* (Maruyama and Kitamura, 1985; Man et al., 1995; Horswill et al., 2001; Starai et al., 2003; Starai and Escalante-Semerena, 2004; Garrity et al., 2007). SCFAs are present at high concentrations in the human gastrointestinal tract (Cummings et al., 1987) and stimulate invasion of the intestinal epithelium by *S. enterica* (Lawhon et al., 2002; Huang et al., 2008; Hung et al., 2013).

Analogously, the SCFAs formate, acetate, propionate and butyrate are among the metabolic signals that elicit differentiation of replicative *L. pneumophila* to the transmissive form, as revealed by phenotypic microarray analysis (Edwards et al., 2009). At concentrations of 10 mM, SCFAs halt *L. pneumophila* growth without affecting cell viability and induce expression of motility, cytotoxicity, sodium sensitivity and avoidance of lysosomal degradation, all hallmark traits of transmissive *L. pneumophila*. The LetA/LetS two-component system is pivotal to the response of *L. pneumophila* to SCFAs as metabolic cues, as *letA* or *letS* mutants are blind to SCFA signals for differentiation. The morphogenetic alterations observed are not merely due to changes in pH, as inorganic acids like hydrochloric acid, even in excess of 10 mM, fail to elicit differentiation of *L. pneumophila* *in vitro*.

A formal possibility was that, to coordinate the response to SCFAs, *L. pneumophila* employs the acetate kinase (Ack)/phosphotransacetylase (Pta) system that converts SCFAs like acetate into their acyl-phosphate (P) or acyl-coenzyme A (CoA) derivatives. This system is utilized by bacteria like *E. coli*, *S. enterica* and *L. monocytogenes* to control several cellular processes (Starai and Escalante-Semerena, 2004; Wolfe, 2005; Gueriri et al., 2008). In this two-enzyme system, acetate may be phosphorylated by Ack with subsequent conversion to acetyl-CoA by Pta. The reactions catalyzed by Ack or Pta can be reversed according to cell needs. The Ack and Pta enzymes may be used by bacteria in several ways: (i) to generate acetyl-P, a high energy intermediate that may serve as a signal for two-component systems involved in activities like biofilm formation, flagellar biosynthesis, motility and pathogenesis (Wolfe et al., 2003; Wolfe, 2005, 2010; Gueriri et al., 2008; Sze and Li, 2011), (ii) to assimilate acetate into acetyl-CoA, or (iii) to dissimilate acetyl-CoA into acetate (Starai and Escalante-Semerena, 2004; Wolfe, 2005).

Unlike in other systems, *L. pneumophila* does not appear to assimilate an excess of these SCFAs through the proteins encoded by *ackA2* and *pta* to relay metabolic signals to the LetA/LetS two-component system. In particular, *L. pneumophila* strains mutated in the genes proposed to encode Ack and Pta nevertheless differentiate from the replicative to the transmissive form in the presence of excess acetate or propionate (Edwards et al., 2009).

Nonetheless, SCFAs do provoke the regulatory cascade that leads to differentiation of replicative *L. pneumophila* to the transmissive form using the stringent response via the bifunctional ppGpp synthetase/hydrolase SpoT enzyme (Dalebroux et al., 2009). It cannot be ruled out, however, that apart from the stringent response, *L. pneumophila* may use pathways distinct to those described for other intracellular pathogens to respond to alterations in fatty acid metabolism and maintain CoA homeostasis when exposed to excess SCFAs.

An additional feature of the integrated nature of metabolism and differentiation in *L. pneumophila* is that the activities of the RelA and SpoT enzymes are dedicated to respond to distinct metabolic signals. Under conditions of fatty acid metabolic stress, ppGpp is synthesized by the bifunctional ppGpp synthetase/hydrolase (SpoT) enzyme. Conversely, RelA is employed by *L. pneumophila* to mount a response to amino acid starvation signals (Dalebroux et al., 2009; Edwards et al., 2009). Intracellular *L. pneumophila* also exhibit distinct requirements for RelA and SpoT as the ppGpp synthetase activity encoded by *relA* is dispensable for intracellular replication, but the ppGpp synthetase/hydrolase activities encoded by the *spoT* gene are not (Dalebroux et al., 2009).

Whereas ppGpp synthesis is required for *L. pneumophila* to express the virulence factors that are required for infection and establishment of the compartment in which it will replicate, ppGpp hydrolysis by SpoT is required for intracellular bacteria to initiate replication (Dalebroux et al., 2009). In fact, *L. pneumophila relA spoT* mutant strains, blocked in ppGpp synthesis, exhibit poor infectivity to macrophages and the bacteria that survive after initial infection are degraded as they are unable to avoid phagosome-lysosome fusion. Experiments in which ppGpp synthetase activity was supplied to *relA spoT* mutant strains by providing *relA*<sup>+</sup> or *spoT*<sup>+</sup> alleles in multicopy indicated that the ppGpp required for infection of the host may be synthesized by either RelA or SpoT. However, the ppGpp synthetase activity of SpoT is specifically required for secondary infections. These data established that *L. pneumophila* requires careful regulation of ppGpp synthesis and hydrolysis to respond to metabolic signals and differentiate accordingly.

When exposed to excess SCFAs, *L. pneumophila relA spoT* mutant strains fail to be cytotoxic to macrophages. Furthermore, *L. pneumophila* strains carrying a particular allele of the *spoT* gene (*spoT-A413E*) in multicopy were blind to perturbations in fatty acid metabolism as they exhibited enhanced cytotoxicity regardless of SCFA exposure (Dalebroux et al., 2009). It was postulated that the increased cytotoxicity phenotype was due to two key features of the mutant SpoT enzyme: increased ppGpp synthetase activity and inability to interact with the fatty acid metabolism machinery of the cell. This line of experiments further demonstrated that the balance between ppGpp synthesis and hydrolysis is critical to the *L. pneumophila* intracellular lifestyle.

In *E. coli*, SpoT activity is modulated via direct interactions with the acyl-carrier protein (ACP) (Battesti and Bouveret, 2006; Angelini et al., 2012), and molecular genetic analyses indicate this is also the case for *L. pneumophila* (Dalebroux et al., 2009). ACP is a crucial component of the fatty acid and phospholipid biosynthetic machinery in bacteria and eukaryotes (Lombard et al.,

2012) that is responsible for carrying fatty acid chains as acyl derivatives attached to a 4'-phosphopantetheine prosthetic group in ACP. Indeed, the interaction of SpoT with ACP requires that the phosphopantetheine moiety be attached to ACP, indicating that SpoT interacts with functional ACP (Battesti and Bouveret, 2006). The structure of ACP is altered by perturbations in fatty acid metabolism, such as inhibition of fatty acid synthesis caused by mutations in fatty acid biosynthetic enzymes (Jackowski and Rock, 1983, 1987). Thus, the conformation of ACP reflects the status of fatty acid metabolism, which it then relays to the SpoT enzyme. SpoT in turn responds by either hydrolyzing ppGpp when fatty acid biosynthesis is uncompromised or synthesizing the alarmone when fatty acid metabolism is perturbed.

*L. pneumophila* also appears to integrate fatty acid metabolism with the expression of virulence factors through alterations in acyl-ACP forms (Edwards et al., 2009). The *L. pneumophila* Philadelphia-1 genome contains several orthologs of genes encoding acetyl-CoA carboxylase (ACC),  $\beta$ -ketoacyl-ACP synthases I (FabB), and II (FabF) (Edwards et al., 2009). When *L. pneumophila* is exposed to known inhibitors of the above mentioned gene products, individually, as in the case of the antibiotic cerulenin, or in combination, it differentiates to the transmissive form. Differentiation under these conditions is proposed to occur by accumulation of fatty acid metabolites that are purported to stall fatty acid metabolism and alter the conformation of ACP. Exposure to SCFAs may also mimic this metabolic state. Analysis of acyl-ACP species of *L. pneumophila* cells that have been exposed to SCFAs lend support to this idea: the acyl-ACP profiles of exponentially replicating bacteria exposed to acetate or propionate mimic those in of cells in the transmissive state (Edwards et al., 2009).

Regardless of whether fatty acid metabolism is altered by pharmacological inhibition of enzyme activity or the presence of particular fatty acid metabolites, both the stringent response commanded by the SpoT enzyme and the LetA/LetS two-component system are required components of the multilayered regulatory strategy employed by *L. pneumophila* to monitor the state of fatty acid metabolism (Dalebroux et al., 2009; Edwards et al., 2009).

## NICOTINIC ACID

The metabolism of NAD<sup>+</sup> and NADP<sup>+</sup> is necessary for redox reactions of catabolism and anabolism in living organisms. In addition, NAD<sup>+</sup> is a required substrate in reactions of DNA synthesis, posttranslational modification of proteins, and cell signaling processes (reviewed in Gazzaniga et al., 2009). The pyridine derivative nicotinic acid is a known component of NAD<sup>+</sup> salvage pathways. Nicotinic acid may be produced from NAD<sup>+</sup> degradation in cells and can be recycled back to NAD<sup>+</sup> via the pyridine nucleotide cycle salvage pathway, also known as the Preiss-Handler pathway (Foster et al., 1979; Foster and Baskowsky-Foster, 1980; Hillyard et al., 1981; Gazzaniga et al., 2009; Houtkooper et al., 2010; Keseler et al., 2011). Exogenous nicotinic acid can also be assimilated via the same route.

Whereas bacteria like *E. coli* and *S. enterica* are capable of *de novo* NAD<sup>+</sup> biosynthesis, microbes like *Bordetella pertussis* and *Candida glabrata* are not (Domergue et al., 2005; Ma et al.,

2007; Gazzaniga et al., 2009). Moreover, the latter organisms can salvage nicotinic acid to satisfy their NAD<sup>+</sup> requirement (McPheat et al., 1983; Domergue et al., 2005; Ma et al., 2007). *B. pertussis* and *C. glabrata* are human pathogens in which nicotinic acid metabolism also regulates the expression of virulence factors such as adhesins, toxins, and agglutinating factors (Schneider and Parker, 1982; MCPheat et al., 1983; Cotter and DiRita, 2000; Domergue et al., 2005).

In *B. pertussis*, nicotinic acid modulates the activity of the BvgA/BvgS two-component system, a master controller of a spectrum of phenotypic states of *B. pertussis* known as Bvg phases (Melton and Weiss, 1989; Cummings et al., 2006). For example, during the Bvg<sup>+</sup> phase, *B. pertussis* is equipped to colonize the human respiratory tract, whereas in the intermediate Bvg<sup>i</sup> phase, *B. pertussis* is more suited for transmission (Vergara-Irigaray et al., 2005). The BvgA/BvgS two-component system works through a four-step phosphorelay that is singular among two-component systems, not only due to the additional steps of the phosphorelay, but also because three of the four phosphorelay domains are found within the BvgS protein itself rather than in three different proteins, as in the *Bacillus subtilis* sporulation system (Appleby et al., 1996). By mechanisms yet unknown, nicotinic acid inactivates the BvgA/BvgS two-component system, thus repressing expression of virulence factors (Schneider and Parker, 1982; MCPheat et al., 1983; Cotter and DiRita, 2000; Cummings et al., 2006). The BvgA/S systems controls panels of traits whose expression depends on the concentration of BvgA-phosphate (Bvg~P), thereby generating a variety of phenotypic states for *B. pertussis* bacteria (Williams and Cotter, 2007; Boulanger et al., 2013).

The LetA/LetS two-component system in *L. pneumophila* has homology to the *B. pertussis* BvgA/BvgS system (Edwards et al., 2010), particularly in the polydomain structure of the LetS protein. The sequence similarities of the two systems and the fact that the LetA/LetS system controls a panel of transmissive traits in *L. pneumophila* led to the idea that nicotinic acid might also modulate gene expression in this bacterium.

A high concentration of nicotinic acid (5 mM) halts growth of replicative *L. pneumophila* and triggers premature expression of transmissive traits, such as motility, evasion of lysosomal degradation, sodium sensitivity, and cytotoxicity to macrophages (Edwards et al., 2013). The LetA/LetS system is required for full induction of the flagellin promoter, but is not required for nicotinic acid to exert its inhibitory effect on growth.

Global transcriptional analyses of replicative *L. pneumophila* exposed to nicotinic acid revealed that this metabolite triggers expression of genes typical of the transmissive state of the bacterium, including those encoding: (i) virulence factors like the *ralF* gene and components of the Dot/Icm apparatus; (ii) the flagellar regulon, and (iii) alternative sigma factors, like *rpoN*, *rpoH*, and *fliA*. Moreover, the transcriptome analyses identified a locus whose expression is altered as a specific response to nicotinic acid exposure. In particular, open-reading frames (ORFs) *lpg0272* and *lpg0273* were induced 9-fold and 35-fold by treatment with nicotinic acid. ORF *lpg0272* is annotated as a cysteine transferase activity, and ORF *lpg0273* is predicted to encode a protein of the major facilitator superfamily (MFS) of transporters. In particular, the predicted protein structure of *Lpg0273* resembles the EmrD

multidrug efflux pump of *E. coli* (Yin et al., 2006; Edwards et al., 2013). Interestingly, two ORFs whose products are proposed to be involved in NAD<sup>+</sup> salvage are encoded immediately upstream of the *lpg0272-3* locus (Caspi et al., 2012; Edwards et al., 2013). Nevertheless, their expression levels were unchanged in response to nicotinic acid (Edwards et al., 2013).

Additional genetic analyses revealed that the product of ORF *lpg0273* equips *L. pneumophila* to cope with the stress of exposure to high concentrations of nicotinic acid. *L. pneumophila* strains that harbor a deletion of the *lpg0273* ORF exhibit increased sensitivity to nicotinic acid. Conversely, strains carrying *lpg0273* in multicopy are more resistance to nicotinic acid (Edwards et al., 2013). Given the properties conferred by *lpg0273* to *L. pneumophila* exposed to nicotinic acid, ORF was designated *mnrA* for “MFS nicotinic acid responder” (Edwards et al., 2013).

The specific mechanisms by which nicotinic acid impacts metabolism and virulence gene expression in *L. pneumophila* and other bacteria still remain to be determined. The transcriptional profiles of transmissive bacteria and replicative bacteria exposed to nicotinic acid harbor several clues that promise to unveil the molecular and metabolic response of *L. pneumophila* to the pyridine compound. First, the loci that displayed the greatest transcriptional changes in response to nicotinic acid are near loci that are proposed to encode NAD<sup>+</sup> salvage functions. Second, one of the ORFs that is highly expressed under nicotinic acid stress confers *L. pneumophila* with protection from this metabolite. This transporter may prove analogous to transporters of *Saccharomyces cerevisiae* that balance import and export of NAD<sup>+</sup> precursors like nicotinamide riboside and nicotinic acid to support NAD<sup>+</sup> homeostasis (Belenky et al., 2011). Third, ORF *lpg0274*, proposed to encode a transcriptional regulator of the LysR family, is positioned 3' of ORF *lpg0273* and is also highly transcribed in the presence of nicotinic acid. This ORF may prove to be a regulator of nicotinic acid metabolism and of the nearby NAD<sup>+</sup> salvage loci in *L. pneumophila*. The coordinated response of *L. pneumophila* to metabolites like nicotinic acid is another display of the strategies this microbe employs to respond to metabolic changes.

## OTHER METABOLITES

Several other metabolites were discovered to trigger premature differentiation of *L. pneumophila*, including, but not limited to, deoxyadenosine, deoxyribose, hydroxylamine, and 4-hydroxybenzoic acid (Edwards et al., 2009). Whether these compounds exert their effect by impacting additional metabolic routes requires additional genetic and biochemical analyses.

## BACTERIAL METABOLIC COMPONENTS REQUIRED FOR VIRULENCE AND INTRACELLULAR GROWTH

### PHAGOSOMAL TRANSPORTERS (Phts) AS NUTRIENT SALVAGING SYSTEMS FOR INTRACELLULAR *L. pneumophila*

The *L. pneumophila* phagosomal (Pht) transporter sub-family of the major facilitator superfamily (MFS) of transporters in Chen et al. (2008) was uncovered through genetic characterization of the *phtA* locus, which is adjacent to *letE*, a locus that encodes an enhancer protein of transmissive trait expression (Bachman and Swanson, 2004). *L. pneumophila* *phtA* mutants exhibit a marked



intracellular growth defect (Sauer et al., 2005). An interesting feature of *phtA* mutants is that they efficiently infect host cells and are able to establish an ER-derived compartment like wild-type *L. pneumophila* (Sauer et al., 2005). However, instead of multiplying like wild-type bacteria, *phtA* mutants remain as single cells in the vacuolar compartment and continue to express transmissive traits (Sauer et al., 2005). Thus, the intracellular growth defect of *phtA* mutants is due to the inability of the bacterial cells to differentiate from the transmissive to the replicative form. The differentiation phenotype of *phtA* mutants can be corrected by addition of threonine, threonine-containing peptides or amino acid supplements, or by supplying the wild-type *phtA* allele *in trans* (Sauer et al., 2005). Given the homology of *phtA* to genes encoding MFS transporters, it was deduced that PhtA affords a threonine acquisition function that equips *L. pneumophila* to differentiate and replicate in macrophages.

Additional molecular, genetic and *in silico* analyses revealed the presence of *phtA* homologues within the *L. pneumophila* Philadelphia-1 genome and within intracellular microbes like *Francisella tularensis* and *Coxiella burnetii* (Sauer et al., 2005; Chen et al., 2008). However, the *L. pneumophila* genome encodes the highest number of Pht functions (Chen et al., 2008).

In addition to *phtA*, 5 additional *pht* loci have been identified as necessary for intracellular replication, namely, *phtC*, *phtD*, *phtE*, *phtF*, and *phtJ* (Sauer, 2006; Fonseca et al., 2014). Whereas, the *phtJ* gene product is required for valine acquisition (Sauer, 2006), the products of the *phtC-D* locus are involved in thymidine salvage in *L. pneumophila* (Fonseca et al., 2014). Thus, *pht* loci are devoted not only to the assimilation of amino acids, but also to additional metabolic functions. The role of the *phtE* and *phtF* loci which are adjacent in the *L. pneumophila* genome remains unknown.

The link of the *phtC-D* locus to thymidine salvage in *L. pneumophila* was deduced by analysis of the *L. pneumophila* genome and phenotypic studies that have been recently reported (Fonseca et al., 2014). The *phtC* and *phtD* genes are located in tandem in the *L. pneumophila* genome with the *tdk* gene, proposed to encode thymidine kinase and immediately upstream of *phtC*, and the *deoB* gene proposed to encode phosphopentomutase and immediately downstream of *phtD*. Since Tdk and DeoB proteins are well-documented thymidine salvage functions in bacteria like *E. coli*, the potential involvement of the *L. pneumophila phtC-D* locus in thymidine salvage was investigated. To assess the role of the *phtC* and *D* gene products in thymidine salvage, we exploited the thymidine auxotrophy of the *L. pneumophila* Lp02 parent strain, conferred by a mutation in *thyA*, encoding the putative thymidylate (dTMP) synthetase enzyme (Berger and Isberg, 1993). Indeed, *phtC* and *phtD* mutants exhibited distinct, but sharp phenotypes when exposed to thymidine starvation conditions *in vitro* (Fonseca et al., 2014). When deprived of exogenous thymidine, *L. pneumophila thyA* strains are highly resilient to starvation conditions, a trait that is surpassed by *thyA phtD* mutant strains. In stark contrast, *thyA phtC* mutant strains not only fail to grow, but quickly lose viability. The thymidine starvation phenotypes of the *thyA phtC* mutant strains are corrected by addition of excess exogenous thymidine or by a *phtC*<sup>+</sup> allele in multicopy.

The sequence homology of PhtC to that of MFS transporters suggests that this protein may transport exogenous thymidine. In fact, *L. pneumophila phtC*<sup>+</sup> in multicopy supplies pyrimidine uptake to an *E. coli* strain that lacks all nucleoside transport systems known for that organism, but *phtD*<sup>+</sup> does not (Fonseca et al., 2014).

The role of the *phtC-D* locus in assimilation of exogenous thymidine in *L. pneumophila* was further tested in isogenic *thyA*<sup>+</sup> strains. The ability of the pyrimidine nucleoside analog 5-fluorodeoxyuridine (FUDR) to block *de novo* dTMP synthesis and thus cell growth was exploited in phenotypic assays of *thyA*<sup>+</sup> *phtC* and *thyA*<sup>+</sup> *phtD* strains. It was initially hypothesized that *phtC* or *phtD* mutant strains, lacking nucleoside uptake functions, would be resistant to growth inhibition by the analog. Surprisingly, the *L. pneumophila thyA*<sup>+</sup> parent is highly resilient to FUDR inhibition, but sensitivity to the analog is increased by *phtC* or *phtD* mutations. In particular, a *phtC* mutation results in full inhibition of growth by FUDR. Again, the phenotypes observed are striking, as impairment of nucleoside transport would result in decreased ability of the analog to enter the cell. FudR inhibition of *L. pneumophila thyA*<sup>+</sup> *phtC* and *thyA*<sup>+</sup> *phtD* strains is corrected by addition of exogenous thymidine and/or *phtC*<sup>+</sup> in multicopy. These results led to the notion that the products of the *phtC-D* locus affect dTMP synthesis by mechanisms that extend beyond nucleoside salvage and that impact *de novo* routes in as yet undefined ways.

The *thyA phtC* and *thyA phtD* growth defects in macrophages were not correctible by addition of exogenous thymidine, restoration of thymidine prototrophy to the mutant strains, or addition of each loci in various combinations (Fonseca et al., 2014). Instead, the whole *tdk-phtC-phtD-deoB* locus was required to complement the intracellular growth defect in the macrophage host. Consistent with this result, mRNA analysis of wild-type *L. pneumophila* grown in the presence of thymidine reveals that under these conditions, *tdk*, *phtC*, *phtD*, and *deoB* are co-transcribed. Thus, whereas under defined *in vitro* conditions *phtC*<sup>+</sup> in multicopy or exogenous thymidine are sufficient to correct the defects of mutants in the *phtC-D* locus, intracellular growth demands more stringent conditions. The distinct and sometimes contrasting thymidine salvage phenotypes exhibited by *phtC* and *phtD* mutant strains suggest that nucleoside metabolism, like in other bacteria, is subject to complex transcriptional and metabolic regulation. Additional molecular and biochemical analyses are required to ascertain the mechanisms by which the *phtC-D* locus contributes to thymidine salvage in *L. pneumophila* and to its intracellular life cycle.

## IRON METABOLISM AND CYTOCHROMES

Like for most living organisms, iron is an essential nutrient for *L. pneumophila* (Reeves et al., 1981). Iron is a constituent of prosthetic groups like heme and iron-sulfur clusters used by several enzymes with diverse and central roles in metabolism such as electron transfer reactions, substrate binding and catalysis, and oxygen transport. The major iron-containing protein in *L. pneumophila* is the homolog to *E. coli* aconitase enzyme of the tricarboxylic acid cycle (Mengaud and Horwitz, 1993). In addition, the genomes of several *L. pneumophila* strains contain genes

proposed to encode iron-sulfur proteins and heme-containing cytochrome *b* and *c*-type cytochromes (Cazalet et al., 2004; Chien et al., 2004; D'Auria et al., 2010; Caspi et al., 2012).

Iron also dictates the outcome of *L. pneumophila* infection. Conditions that increase iron availability, like iron supplementation of tissue cultures or feeding of iron to animal hosts, favor *L. pneumophila* infection (reviewed in Cianciotto, 2007), whereas limiting iron availability attenuates replication (James et al., 1995; Byrd and Horwitz, 2000; Viswanathan et al., 2000).

*L. pneumophila* possesses several systems to salvage and assimilate iron from its environment: the siderophore legiobactin (Liles et al., 2000; Allard et al., 2006), ferric reductase enzymes (Johnson et al., 1991; Poch and Johnson, 1993; James et al., 1997), the FeoB ferrous iron transporter (Robey and Cianciotto, 2002), the *iraAB* locus (Pope et al., 1996; Viswanathan et al., 2000) and two secreted ferric iron reductants, homogenistic acid, and homogenistic acid-melanin (pyomelanin) (Zheng et al., 2013). In addition, both the cytochrome *c* maturation (Ccm) system and cytochrome *c*<sub>4</sub> are required for expression of the legiobactin siderophore of *L. pneumophila* (Yip et al., 2011). The specific cellular function of Ccm system proteins is to attach heme covalently to cytochrome *c* precursor proteins. The link between Ccm function and iron salvage and assimilation is not unprecedented, since other bacterial species also display this type of metabolic integration (Cianciotto et al., 2005).

The phenotypes of mutants defective in some of the iron salvaging and utilization systems highlight the importance of iron for intracellular replication of *L. pneumophila*. First, mutants defective in iron assimilation (e.g., *feoB*, *iraA*) exhibit reduced intracellular replication, a defect exacerbated when host cells are pre-treated with iron chelators (Viswanathan et al., 2000; Robey and Cianciotto, 2002). Second, mutants of the *frgA* gene, proposed to encode a homolog of *E. coli* aerobactin siderophore synthetases, are impaired in replication in macrophages. These data indicate a requirement for at least one siderophore during intracellular growth, as legiobactin is dispensable under these conditions (Hickey and Cianciotto, 1997; Allard et al., 2006; Cianciotto, 2007). Third, mutants defective in cytochrome *c*-maturation (*ccm*) exhibit replication defects in macrophages and amoeba (Viswanathan et al., 2002; Yip et al., 2011) and in a mouse pulmonary model of infection (Naylor and Cianciotto, 2004). The defects of *ccm* mutants in macrophages are alleviated by addition of iron and exacerbated by iron depletion (Viswanathan et al., 2002; Naylor and Cianciotto, 2004). Furthermore, studies of *c*-type cytochrome mutants indicate that *L. pneumophila* requires cytochromes *c*<sub>5</sub> and *c*<sub>1</sub> to grow in macrophages or amoeba, indicating that poor growth by *ccm* mutants may reflect an inability of *L. pneumophila* to generate the aforementioned cytochromes (Yip et al., 2011). Given the pivotal contributions of *c*-type cytochromes in aerobic respiration, it is easy to surmise why *L. pneumophila* requires these proteins for intracellular growth. Whether *c*-type cytochromes have additional roles during intracellular growth of *L. pneumophila* is an intriguing possibility that is worthy of consideration in light of various additional roles for these proteins discovered in other systems (Yip et al., 2011).

## HOST METABOLIC COMPONENTS EXPLOITED BY INTRACELLULAR *L. pneumophila*

### HOST AMINO ACID TRANSPORTERS

The notion that *L. pneumophila* absolutely relies on certain host metabolic functions for intracellular replication is demonstrated by an elegant study by Wieland et al. (2005), whereby SLC1A5, a Na<sup>+</sup>/Cl<sup>-</sup> coupled amino acid transporter of eukaryotes, was identified as necessary for *L. pneumophila* replication in MM6-monocytes. Furthermore, replication by intracellular bacteria required that MM6 cell culture medium be supplemented with amino acids. Analyses of host mRNAs indicated that the SLC1A5 transporter is highly expressed in infected MM6 monocytes. Moreover, pharmacological and siRNA inhibition of the SLC1A5 transporter blocks *L. pneumophila* replication, providing strong evidence that the host amino acid transporter is required for bacterial growth. Thus, by exploiting host amino acid acquisition systems, *L. pneumophila* fuels its own carbon and energy metabolism.

This study also uncovered some differences between the amino acid requirements of intracellular and extracellular bacteria. For instance, L-glutamine supports optimal growth in host cells and in broth cultures (Wieland et al., 2005). Conversely, L-glutamate, L-tyrosine and L-threonine are required for growth in broth cultures but have minimal impact on intracellular growth. From studies of this type, it is possible to predict classes of metabolites that are readily available within host cells.

### THE HOST PROTEASOME

*L. pneumophila* utilizes the Dot/Icm apparatus to translocate a series of effectors across the bacterial plasma membrane. One of these effectors is the AnkB protein (Al-Khodori et al., 2008; Price et al., 2009, 2011; Al-Quadan and Kwai, 2011; Bruckert et al., 2014). The AnkB protein contains several eukaryotic domains, including an F-box domain (Price et al., 2009; Skaar et al., 2013), which has a well-documented role in ubiquitination. This type of posttranslational modification regulates a variety of cellular processes, such as protein degradation, cell division, and protein trafficking (Schnell and Hicke, 2003; Hershko, 2005; Price et al., 2009; Komander and Rape, 2012).

*L. pneumophila* require AnkB to replicate in various hosts, including U937 macrophages, *A. castellanii*, *A. polyphaga*, as well as for pulmonary infection in a mouse model (Al-Khodori et al., 2008; Lomma et al., 2010; Price et al., 2010). The AnkB effector itself is posttranslationally modified by farnesylation, which mediates its localization on the membrane of the *L. pneumophila* replication vacuole (Price et al., 2010). *L. pneumophila* fails to replicate intracellularly when farnesylation of AnkB is blocked, highlighting the importance of farnesylation and the requirement for the effector in host cells (Price et al., 2009; Al-Quadan and Kwai, 2011). Anchoring of AnkB to the vacuolar membrane enables the effector to serve as a dock for polyubiquitinated proteins, which are then degraded by the host proteasome (Price et al., 2011). Degradation of polyubiquitinated proteins was deduced to generate a pool of amino acids that support *L. pneumophila* replication within the host (Price et al., 2011). After infection, *L. pneumophila* *ankB* mutants prematurely differentiate to the transmissive form, as judged by expression of



*flaA*, *relA* and *spoT*, three genes that are induced by starvation *in vitro*. However, when the infected cells are supplemented with excess amino acids, *ankB* mutants replicate intracellularly and do not activate the *flaA* promoter. The tight link between host metabolism and the *L. pneumophila* life cycle is further evidenced by the finding that swift farnesylation of AnkB and subsequent assembly and degradation of polyubiquitinated proteins occurs upon attachment of *L. pneumophila* to host cells (Bruckert et al., 2014). This in turn is thought to increase amino acid levels in the host to promote the transition of *L. pneumophila* from the transmissive to the replicative phase of its intracellular life cycle.

### HOST PHOSPHOLIPIDS

The finding that *L. pneumophila* couples fatty acid metabolism to virulence strongly indicated that host lipid or fatty acid metabolites may serve both as nutrients and as signals that relay the metabolic state of the host and the bacterium (Dalebroux et al., 2009; Edwards et al., 2009). Further strengthening this notion is the fact that *L. pneumophila* secretes a number of phospholipase enzymes, some of which have been determined to be virulence factors that act by destroying host membranes (Flieger et al., 2000, 2002, 2004; Bender et al., 2009; Kuhle and Flieger, 2013).

In addition, *L. pneumophila* also synthesizes several effectors that are translocated by the Dot/Icm type IV secretion system that act on phospholipids and modify the phagosomal compartment in which the microbe replicates (VanRheenen et al., 2006; Viner et al., 2012). While much of this work has focused on identification of effectors and their activities, a hypothesis that warrants experimental testing is that *L. pneumophila* deploys effectors not only to alter organelle membranes and signaling pathways in its host, but also to salvage lipid precursors.

### POLYAMINES

Polyamines, such as spermidine and putrescine, are well known for their effects on cell growth, nucleic acid synthesis, and cell integrity in prokaryotic and eukaryotic organisms (reviewed in Tabor and Tabor, 1985; Pegg, 1986; Chattopadhyay and Tabor, 2013). However, the detailed mechanisms by which these polycationic compounds affect various cellular processes have been the object of ongoing study for several decades and are still being determined.

Microorganisms like *E. coli* and *S. cerevisiae* have served as amenable models for defining the effects of polyamines on cell physiology (Hafner et al., 1979; Geiger and Morris, 1980; Abraham, 1981; Tabor and Tabor, 1985; Chattopadhyay et al., 2009a,b; Chattopadhyay and Tabor, 2013). In *E. coli*, polyamines contribute to optimal aerobic growth, synthesis of proteins and nucleic acids, protection from oxidative stress, and anaerobic growth (Tabor et al., 1980; Chattopadhyay et al., 2009a). A capacity for polyamines to protect *E. coli* from acid stress was recently discovered (Chattopadhyay and Tabor, 2013).


Polyamines have been shown to influence various aspects of microbial pathogenesis (reviewed in Shah and Swiatlo, 2008). Of particular interest is that microbes like *Helicobacter pylori* and *Pneumocystis jirovecii* perturb metabolism of polyamines in host cells (Lasbury et al., 2007; Chaturvedi et al., 2012a,b; Xu et al., 2012). The alterations in polyamine metabolism caused by these

microbes are thought to trigger apoptosis in the case of *P. jirovecii* and to be linked to *H. pylori*-associated gastric diseases such as cancer.

Polyamines also appear to contribute to replication of intracellular *L. pneumophila*. *L. pneumophila* encodes an HtpB chaperonin (Nasrallah et al., 2011), a protein known in some systems to recruit host mitochondria and alter microfilament arrangements (Chong et al., 2009). In a yeast two-hybrid system, HtpB interacts with eukaryotic S-adenosylmethionine decarboxylase (SAMDC), an enzyme prokaryotes and eukaryotes require for synthesis of spermidine (Caspi et al., 2012). Furthermore, when SAMDC activity is inhibited in U937 macrophages infected with *L. pneumophila*, intracellular replication of the bacterium is significantly reduced (Nasrallah et al., 2011). Given that *L. pneumophila* appears to lack most canonical enzymes required for polyamine synthesis, it has been proposed that *L. pneumophila* salvages any polyamines that promote its replication (Nasrallah et al., 2011). Indeed, exogenous polyamines boost growth of intracellular and extracellular *L. pneumophila*. However, it has not been ruled out that, like other microbial pathogens, *L. pneumophila* may manipulate SAMDC activity to alter the physiology of the host to its advantage. Further studies of the links between host polyamine metabolism and *L. pneumophila* infection promise to increase our understanding of how specific host systems are exploited for microbial pathogenesis.

### CONCLUDING REMARKS

For a number of years, there has been considerable emphasis on understanding the mechanisms of pathogenesis of intracellular microbes like *L. pneumophila*. Two key aspects of *L. pneumophila*



virulence have been object of intense study: (i) the strategies employed by the bacterium to manipulate host cellular processes and (ii) the coupling of its cellular differentiation cycle with virulence. Invariably, both lines of research have discovered intricate connections between microbial metabolism and mechanisms of disease. In several examples presented herein, whether *L. pneumophila* replicates or deploys virulence factors to seek a new host is dictated by the presence of particular metabolites, such as amino acids or fatty acids. In some instances the source of these metabolites is clear (i.e., the host cell), in others, it remains an open question (Figure 2). It is also becoming increasingly clear that at least some of the effectors produced by *L. pneumophila* upon infection are geared toward exploitation of the host as a nutritional niche (Figure 2).

*In vitro* studies have been instrumental in gaining insights to the nutritional requirements and metabolic capabilities of *L. pneumophila*. These together with the experimental approaches taken to dissect the *L. pneumophila* cell differentiation cycle in synchronous broth cultures have laid the foundation to explore further the metabolic strategies employed by *L. pneumophila* to thrive intracellularly. Emerging genetic, molecular and biochemical methodologies increase the number of tools that are available to answer the many questions that remain to be answered regarding the metabolism of *L. pneumophila* in host cells.

## REFERENCES

- Abdel-Nour, M., Duncan, C., Low, D. E., and Guyard, C. (2013). Biofilms: the stronghold of *Legionella pneumophila*. *Int. J. Mol. Sci.* 14, 21660–21675. doi: 10.3390/ijms141121660
- Abraham, A. K. (1981). Effect of polyamines on the fidelity of macromolecular synthesis. *Med. Biol.* 59, 368–373.
- Al-Bana, B. H., Haddad, M. T., and Garduno, R. A. (2014). Stationary phase and mature infectious forms of *Legionella pneumophila* produce distinct viable but non-culturable cells. *Environ. Microbiol.* 16, 382–395. doi: 10.1111/1462-2920.12219
- Albert-Weissenberger, C., Sahr, T., Sismeiro, O., Hacker, J., Heuner, K., and Buchrieser, C. (2010). Control of flagellar gene regulation in *Legionella pneumophila* and its relation to growth phase. *J. Bacteriol.* 192, 446–455. doi: 10.1128/JB.00610-09
- Al-Khodori, S., Price, C. T., Habyarimana, F., Kalia, A., and Abu Kwaik, Y. (2008). A Dot/Icm-translocated ankyrin protein of *Legionella pneumophila* is required for intracellular proliferation within human macrophages and protozoa. *Mol. Microbiol.* 70, 908–923. doi: 10.1111/j.1365-2958.2008.06453.x
- Allard, K. A., Viswanathan, V. K., and Cianciotto, N. P. (2006). IbtA and IbtB are required for production of the *Legionella pneumophila* siderophore legiobactin. *J. Bacteriol.* 188, 1351–1363. doi: 10.1128/JB.188.4.1351-1363.2006
- Allen, S., Zilles, J. L., and Downs, D. M. (2002). Metabolic flux in both the purine mononucleotide and histidine biosynthetic pathways can influence synthesis of the hydroxymethyl pyrimidine moiety of thiamine in *Salmonella enterica*. *J. Bacteriol.* 184, 6130–6137. doi: 10.1128/JB.184.22.6130-6137.2002
- Al-Quadan, T., and Kwaik, Y. A. (2011). Molecular characterization of exploitation of the polyubiquitination and farnesylation machineries of *Dictyostelium discoideum* by the AnkB F-Box effector of *Legionella pneumophila*. *Front. Microbiol.* 2:23. doi: 10.3389/fmicb.2011.00023
- Al-Quadan, T., Price, C. T., and Abu Kwaik, Y. (2012). Exploitation of evolutionarily conserved amoeba and mammalian processes by *Legionella*. *Trends Microbiol.* 20, 299–306. doi: 10.1016/j.tim.2012.03.005
- Amyot, W. M., deJesus, D., and Isberg, R. R. (2013). Poison domains block transit of translocated substrates via the *Legionella pneumophila* Icm/Dot system. *Infect. Immun.* 81, 3239–3252. doi: 10.1128/IAI.00552-13
- Angelini, S., My, L., and Bouveret, E. (2012). Disrupting the Acyl Carrier Protein/SpoT interaction *in vivo*: identification of ACP residues involved in the interaction and consequence on growth. *PLoS ONE* 7:e36111. doi: 10.1371/journal.pone.0036111
- Appleby, J. L., Parkinson, J. S., and Bourret, R. B. (1996). Signal transduction via the multi-step phosphorelay: not necessarily a road less traveled. *Cell* 86, 845–848. doi: 10.1016/S0092-8674(00)80158-0
- Bachman, M. A., and Swanson, M. S. (2001). RpoS co-operates with other factors to induce *Legionella pneumophila* virulence in the stationary phase. *Mol. Microbiol.* 40, 1201–1214. doi: 10.1046/j.1365-2958.2001.02465.x
- Bachman, M. A., and Swanson, M. S. (2004). The LetE protein enhances expression of multiple LetA/LetS-dependent transmission traits by *Legionella pneumophila*. *Infect. Immun.* 72, 3284–3293. doi: 10.1128/IAI.72.6.3284-3293.2004
- Battesti, A., and Bouveret, E. (2006). Acyl carrier protein/SpoT interaction, the switch linking SpoT-dependent stress response to fatty acid metabolism. *Mol. Microbiol.* 62, 1048–1063. doi: 10.1111/j.1365-2958.2006.05442.x
- Bazurto, J. V., and Downs, D. M. (2014). Amino-4-imidazolecarboxamide ribotide (AICAR) directly inhibits coenzyme A biosynthesis in *Salmonella enterica*. *J. Bacteriol.* 196, 772–779. doi: 10.1128/JB.01087-13
- Belenky, P., Stebbins, R., Bogan, K. L., Evans, C. R., and Brenner, C. (2011). NrtI and TnaI-independent export of NAD<sup>+</sup> precursor vitamins promotes NAD<sup>+</sup> homeostasis and allows engineering of vitamin production. *PLoS ONE* 6:e19710. doi: 10.1371/journal.pone.0019710
- Bender, J., Rydzewski, K., Broich, M., Schunder, E., Heuner, K., and Flieger, A. (2009). Phospholipase PlA<sub>B</sub> of *Legionella pneumophila* represents a novel lipase family: protein residues essential for lipolytic activity, substrate specificity, and hemolysis. *J. Biol. Chem.* 284, 27185–27194. doi: 10.1074/jbc.M109.026021
- Berger, K. H., and Isberg, R. R. (1993). Two distinct defects in intracellular growth complemented by a single genetic locus in *Legionella pneumophila*. *Mol. Microbiol.* 7, 7–19. doi: 10.1111/j.1365-2958.1993.tb01092.x
- Boulanger, A., Chen, Q., Hinton, D. M., and Stibitz, S. (2013). *In vivo* phosphorylation dynamics of the *Bordetella pertussis* virulence-controlling response regulator BvgA. *Mol. Microbiol.* 88, 156–172. doi: 10.1111/mmi.12177
- Browning, D. F., and Busby, S. J. (2004). The regulation of bacterial transcription initiation. *Nat. Rev. Microbiol.* 2, 57–65. doi: 10.1038/nrmicro787
- Bruckert, W. M., Price, C. T., and Abu Kwaik, Y. (2014). Rapid nutritional remodeling of the host cell upon attachment of *Legionella pneumophila*. *Infect. Immun.* 82, 72–82. doi: 10.1128/IAI.01079-13
- Bruggemann, H., Hagman, A., Jules, M., Sismeiro, O., Dillies, M. A., Gouyette, C., et al. (2006). Virulence strategies for infecting phagocytes deduced from the *in vivo* transcriptional program of *Legionella pneumophila*. *Cell. Microbiol.* 8, 1228–1240. doi: 10.1111/j.1462-5822.2006.00703.x
- Bryan, A., Harada, K., and Swanson, M. S. (2011). Efficient generation of unmarked deletions in *Legionella pneumophila*. *Appl. Environ. Microbiol.* 77, 2545–2548. doi: 10.1128/AEM.02904-10
- Buchrieser, C., and Charpentier, X. (2013). Induction of competence for natural transformation in *Legionella pneumophila* and exploitation for mutant construction. *Methods Mol. Biol.* 954, 183–195. doi: 10.1007/978-1-62703-161-5\_9
- Byrd, T. F., and Horwitz, M. A. (2000). Aberrantly low transferrin receptor expression on human monocytes is associated with nonpermissiveness for *Legionella pneumophila* growth. *J. Infect. Dis.* 181, 1394–1400. doi: 10.1086/315390
- Byrne, B., and Swanson, M. S. (1998). Expression of *Legionella pneumophila* virulence traits in response to growth conditions. *Infect. Immun.* 66, 3029–3034.
- Caspi, R., Altman, T., Dreher, K., Fulcher, C. A., Subhraveti, P., Keseler, I. M., et al. (2012). The MetaCyc database of metabolic pathways and enzymes and the BioCyc collection of pathway/genome databases. *Nucl. Acids Res.* 40, D742–D753. doi: 10.1093/nar/gkr1014
- Cazalet, C., Rusniok, C., Bruggemann, H., Zidane, N., Magnier, A., Ma, L., et al. (2004). Evidence in the *Legionella pneumophila* genome for exploitation of host cell functions and high genome plasticity. *Nat. Genet.* 36, 1165–1173. doi: 10.1038/ng1447
- Chattopadhyay, M. K., Chen, W., Poy, G., Cam, M., Stiles, D., and Tabor, H. (2009b). Microarray studies on the genes responsive to the addition of spermidine or spermine to a *Saccharomyces cerevisiae* spermidine synthase mutant. *Yeast* 26, 531–544. doi: 10.1002/yea.1703
- Chattopadhyay, M. K., Tabor, C. W., and Tabor, H. (2009a). Polyamines are not required for aerobic growth of *Escherichia coli*: preparation of a strain with deletions in all of the genes for polyamine biosynthesis. *J. Bacteriol.* 191, 5549–5552. doi: 10.1128/JB.00381-09

- Chattopadhyay, M. K., and Tabor, H. (2013). Polyamines are critical for the induction of the glutamate decarboxylase-dependent acid resistance system in *Escherichia coli*. *J. Biol. Chem.* 288, 33559–33570. doi: 10.1074/jbc.M113.510552
- Chaturvedi, R., de Sablet, T., Peek, R. M., and Wilson, K. T. (2012a). Spermine oxidase, a polyamine catabolic enzyme that links *Helicobacter pylori* CagA and gastric cancer risk. *Gut Microbes* 3, 48–56. doi: 10.4161/gmic.19345
- Chaturvedi, R., de Sablet, T., Coburn, L. A., Gobert, A. P., and Wilson, K. T. (2012b). Arginine and polyamines in *Helicobacter pylori*-induced immune dysregulation and gastric carcinogenesis. *Amino Acids* 42, 627–640. doi: 10.1007/s00726-011-1038-4
- Chen, D. E., Podell, S., Sauer, J. D., Swanson, M. S., and Saier, M. H. Jr. (2008). The phagosomal nutrient transporter (Pht) family. *Microbiology* 154, 42–53. doi: 10.1099/mic.0.2007/010611-0
- Chien, M., Morozova, L., Shi, S., Sheng, H., Chen, J., Gomez, S. M., et al. (2004). The genomic sequence of the accidental pathogen *Legionella pneumophila*. *Science* 305, 1966–1968. doi: 10.1126/science.1099776
- Chong, A., Lima, C. A., Allan, D. S., Nasrallah, G. K., and Garduno, R. A. (2009). The purified and recombinant *Legionella pneumophila* chaperonin alters mitochondrial trafficking and microfilament organization. *Infect. Immun.* 77, 4724–4739. doi: 10.1128/IAI.00150-09
- Cianciotto, N. P. (2007). Iron acquisition by *Legionella pneumophila*. *Biomaterials* 20, 323–331. doi: 10.1007/s10534-006-9057-4
- Cianciotto, N. P., Cornelis, P., and Baysse, C. (2005). Impact of the bacterial type I cytochrome c maturation system on different biological processes. *Mol. Microbiol.* 56, 1408–1415. doi: 10.1111/j.1365-2958.2005.04650.x
- Cotter, P. A., and DiRita, V. J. (2000). Bacterial virulence gene regulation: an evolutionary perspective. *Annu. Rev. Microbiol.* 54, 519–565. doi: 10.1146/annurev.micro.54.1.519
- Cummings, C. A., Bootsma, H. J., Relman, D. A., and Miller, J. F. (2006). Species- and strain-specific control of a complex, flexible regulon by Bordetella BvgAS. *J. Bacteriol.* 188, 1775–1785. doi: 10.1128/JB.188.5.1775-1785.2006
- Cummings, J. H., Pomare, E. W., Branch, W. J., Naylor, C. P., and Macfarlane, G. T. (1987). Short chain fatty acids in human large intestine, portal, hepatic and venous blood. *Gut* 28, 1221–1227. doi: 10.1136/gut.28.10.1221
- Dalebroux, Z. D., Edwards, R. L., and Swanson, M. S. (2009). SpoT governs *Legionella pneumophila* differentiation in host macrophages. *Mol. Microbiol.* 71, 640–658. doi: 10.1111/j.1365-2958.2008.06555.x
- Dalebroux, Z. D., Yagi, B. F., Sahr, T., Buchrieser, C., and Swanson, M. S. (2010). Distinct roles of ppGpp and DksA in *Legionella pneumophila* differentiation. *Mol. Microbiol.* 76, 200–219. doi: 10.1111/j.1365-2958.2010.07094.x
- D'Auria, G., Jimenez-Hernandez, N., Peris-Bondia, F., Moya, A., and Latorre, A. (2010). *Legionella pneumophila* pangenome reveals strain-specific virulence factors. *BMC Genomics* 11:181. doi: 10.1186/1471-2164-11-181
- Domergue, R., Castano, I., De Las Penas, A., Zupancic, M., Lockatell, V., Hebel, J. R., et al. (2005). Nicotinic acid limitation regulates silencing of *Candida adhesins* during UTI. *Science* 308, 866–870. doi: 10.1126/science.1108640
- Durfée, T., Hansen, A. M., Zhi, H., Blattner, F. R., and Jin, D. J. (2008). Transcription profiling of the stringent response in *Escherichia coli*. *J. Bacteriol.* 190, 1084–1096. doi: 10.1128/JB.01092-07
- Edwards, R. L., Bryan, A., Jules, M., Harada, K., Buchrieser, C., and Swanson, M. S. (2013). Nicotinic acid modulates *Legionella pneumophila* gene expression and induces virulence traits. *Infect. Immun.* 81, 945–955. doi: 10.1128/IAI.00999-12
- Edwards, R. L., Dalebroux, Z. D., and Swanson, M. S. (2009). *Legionella pneumophila* couples fatty acid flux to microbial differentiation and virulence. *Mol. Microbiol.* 71, 1190–1204. doi: 10.1111/j.1365-2958.2008.06593.x
- Edwards, R. L., Jules, M., Sahr, T., Buchrieser, C., and Swanson, M. S. (2010). The *Legionella pneumophila* LetA/LetS two-component system exhibits rheostat-like behavior. *Infect. Immun.* 78, 2571–2583. doi: 10.1128/IAI.01107-09
- Ensminger, A. W., and Isberg, R. R. (2009). *Legionella pneumophila* Dot/Icm translocated substrates: a sum of parts. *Curr. Opin. Microbiol.* 12, 67–73. doi: 10.1016/j.mib.2008.12.004
- Eylert, E., Herrmann, V., Jules, M., Gillmaier, N., Lautner, M., Buchrieser, C., et al. (2010). Isotopologue profiling of *Legionella pneumophila*: role of serine and glucose as carbon substrates. *J. Biol. Chem.* 285, 22232–22243. doi: 10.1074/jbc.M110.128678
- Faucher, S. P., Mueller, C. A., and Shuman, H. A. (2011). *Legionella pneumophila* transcriptome during intracellular multiplication in human macrophages. *Front. Microbiol.* 2:60. doi: 10.3389/fmicb.2011.00060
- Faulkner, G., Berk, S. G., Garduno, E., Ortiz-Jimenez, M. A., and Garduno, R. A. (2008). Passage through *Tetrahymena tropicalis* triggers a rapid morphological differentiation in *Legionella pneumophila*. *J. Bacteriol.* 190, 7728–7738. doi: 10.1128/JB.00751-08
- Faulkner, G., and Garduno, R. A. (2002). Ultrastructural analysis of differentiation in *Legionella pneumophila*. *J. Bacteriol.* 184, 7025–7041. doi: 10.1128/JB.184.24.7025-7041.2002
- Fernandez-Moreira, E., Helbig, J. H., and Swanson, M. S. (2006). Membrane vesicles shed by *Legionella pneumophila* inhibit fusion of phagosomes with lysosomes. *Infect. Immun.* 74, 3285–3295. doi: 10.1128/IAI.01382-05
- Flieger, A., Gongab, S., Faigle, M., Mayer, H. A., Kehrer, U., Mussotter, J., et al. (2000). Phospholipase A secreted by *Legionella pneumophila* destroys alveolar surfactant phospholipids. *FEMS Microbiol. Lett.* 188, 129–133. doi: 10.1111/j.1574-6968.2000.tb09183.x
- Flieger, A., Neumeister, B., and Cianciotto, N. P. (2002). Characterization of the gene encoding the major secreted lysophospholipase A of *Legionella pneumophila* and its role in detoxification of lysophosphatidylcholine. *Infect. Immun.* 70, 6094–6106. doi: 10.1128/IAI.70.11.6094-6106.2002
- Flieger, A., Rydzewski, K., Banerji, S., Broich, M., and Heuner, K. (2004). Cloning and characterization of the gene encoding the major cell-associated phospholipase A of *Legionella pneumophila*, plaB, exhibiting hemolytic activity. *Infect. Immun.* 72, 2648–2658. doi: 10.1128/IAI.72.5.2648-2658.2004
- Fonseca, M. V., Sauer, J. D., Crepin, S., Byrne, B., and Swanson, M. S. (2014). The phtC-phtD locus equips *Legionella pneumophila* for thymidine salvage and replication in macrophages. *Infect. Immun.* 82, 720–730. doi: 10.1128/IAI.01043-13
- Fonseca, M. V., Sauer, J. D., and Swanson, M. S. (2008). “Nutrient acquisition and assimilation strategies of *Legionella pneumophila*,” in *Legionella Molecular Microbiology*, eds K. Heuner and M. S. Swanson (Norwich: Horizon Scientific Press), 213–226.
- Foster, J. W., and Baskowsky-Foster, A. M. (1980). Pyridine nucleotide cycle of *Salmonella typhimurium*: in vivo recycling of nicotinamide adenine dinucleotide. *J. Bacteriol.* 142, 1032–1035.
- Foster, J. W., Kinney, D. M., and Moat, A. G. (1979). Pyridine nucleotide cycle of *Salmonella typhimurium*: regulation of nicotinic acid phosphoribosyltransferase and nicotinamide deamidase. *J. Bacteriol.* 138, 957–961.
- Garduno, R. A. (2008). “Developmental cycle - differentiation of *Legionella pneumophila*,” in *Legionella Molecular Microbiology*, eds K. Heuner and M. S. Swanson (Norwich: Horizon Scientific Press), 55–73.
- Garduno, R. A., Garduno, E., Hiltz, M., and Hoffman, P. S. (2002). Intracellular growth of *Legionella pneumophila* gives rise to a differentiated form dissimilar to stationary-phase forms. *Infect. Immun.* 70, 6273–6283. doi: 10.1128/IAI.70.11.6273-6283.2002
- Garrity, J., Gardner, J. G., Hawse, W., Wolberger, C., and Escalante-Semerena, J. C. (2007). N-lysine propionylation controls the activity of propionyl-CoA synthetase. *J. Biol. Chem.* 282, 30239–30245. doi: 10.1074/jbc.M704409200
- Gazzaniga, F., Stebbins, R., Chang, S. Z., McPeck, M. A., and Brenner, C. (2009). Microbial NAD metabolism: lessons from comparative genomics. *Microbiol. Mol. Biol. Rev.* 73, 529–541. doi: 10.1128/MMBR.00042-08
- Geiger, L. E., and Morris, D. R. (1980). Stimulation of deoxyribonucleic acid replication fork movement by spermidine analogs in polyamine-deficient *Escherichia coli*. *J. Bacteriol.* 141, 1192–1198.
- Gentry, D. R., Hernandez, V. J., Nguyen, L. H., Jensen, D. B., and Cashel, M. (1993). Synthesis of the stationary-phase sigma factor sigma S is positively regulated by ppGpp. *J. Bacteriol.* 175, 7982–7989.
- George, J. R., Pine, L., Reeves, M. W., and Harrell, W. K. (1980). Amino acid requirements of *Legionella pneumophila*. *J. Clin. Microbiol.* 11, 286–291.
- Greub, G., and Raoult, D. (2003). Morphology of *Legionella pneumophila* according to their location within *Hartmannella vermiformis*. *Res. Microbiol.* 154, 619–621. doi: 10.1016/j.resmic.2003.08.003
- Gueriri, I., Bay, S., Dubrac, S., Cyncynatus, C., and Msadek, T. (2008). The Pta-AckA pathway controlling acetyl phosphate levels and the phosphorylation state of the DegU orphan response regulator both play a role in regulating *Listeria monocytogenes* motility and chemotaxis. *Mol. Microbiol.* 70, 1342–1357. doi: 10.1111/j.1365-2958.2008.06496.x
- Hafner, E. W., Tabor, C. W., and Tabor, H. (1979). Mutants of *Escherichia coli* that do not contain 1,4-diaminobutane (putrescine) or spermidine. *J. Biol. Chem.* 254, 12419–12426.



- Hammer, B. K., and Swanson, M. S. (1999). Co-ordination of *Legionella pneumophila* virulence with entry into stationary phase by ppGpp. *Mol. Microbiol.* 33, 721–731. doi: 10.1046/j.1365-2958.1999.01519.x
- Hammer, B. K., Tateda, E. S., and Swanson, M. S. (2002). A two-component regulator induces the transmission phenotype of stationary-phase *Legionella pneumophila*. *Mol. Microbiol.* 44, 107–118. doi: 10.1046/j.1365-2958.2002.02884.x
- Henkin, T. M. (2000). Transcription termination control in bacteria. *Curr. Opin. Microbiol.* 3, 149–153. doi: 10.1016/S1369-5274(00)00067-9
- Hershko, A. (2005). The ubiquitin system for protein degradation and some of its roles in the control of the cell division cycle. *Cell Death Differ.* 12, 1191–1197. doi: 10.1038/sj.cdd.4401702
- Heuner, K., and Eisenreich, W. (2013). The intracellular metabolism of legionella by isotopologue profiling. *Methods Mol. Biol.* 954, 163–181. doi: 10.1007/978-1-62703-161-5\_8
- Hickey, E. K., and Cianciotto, N. P. (1997). An iron- and fur-repressed *Legionella pneumophila* gene that promotes intracellular infection and encodes a protein with similarity to the *Escherichia coli* aerobactin synthetases. *Infect. Immun.* 65, 133–143.
- Hillyard, D., Rechsteiner, M., Manlapaz-Ramos, P., Imperial, J. S., Cruz, L. J., and Olivera, B. M. (1981). The pyridine nucleotide cycle. Studies in *Escherichia coli* and the human cell line D98/AH2. *J. Biol. Chem.* 256, 8491–8497.
- Horswill, A. R., Dudding, A. R., and Escalante-Semerena, J. C. (2001). Studies of propionate toxicity in *Salmonella enterica* identify 2-methylcitrate as a potent inhibitor of cell growth. *J. Biol. Chem.* 276, 19094–19101. doi: 10.1074/jbc.M100244200
- Houtkooper, R. H., Canto, C., Wanders, R. J., and Auwerx, J. (2010). The secret life of NAD<sup>+</sup>: an old metabolite controlling new metabolic signaling pathways. *Endocr. Rev.* 31, 194–223. doi: 10.1210/er.2009-0026
- Hovel-Miner, G., Faucher, S. P., Charpentier, X., and Shuman, H. A. (2010). ArgR-regulated genes are derepressed in the *Legionella*-containing vacuole. *J. Bacteriol.* 192, 4504–4516. doi: 10.1128/JB.00465-10
- Hovel-Miner, G., Pampou, G., Faucher, S. P., Clarke, M., Morozova, I., Morozov, P., et al. (2009). SigmaS controls multiple pathways associated with intracellular multiplication of *Legionella pneumophila*. *J. Bacteriol.* 191, 2461–2473. doi: 10.1128/JB.01578-08
- Huang, Y., Suyemoto, M., Garner, C. D., Cicconi, K. M., and Altier, C. (2008). Formate acts as a diffusible signal to induce *Salmonella* invasion. *J. Bacteriol.* 190, 4233–4241. doi: 10.1128/JB.00205-08
- Hung, C. C., Garner, C. D., Schlauch, J. M., Dwyer, Z. W., Lawhon, S. D., Frye, J. G., et al. (2013). The intestinal fatty acid propionate inhibits *Salmonella* invasion through the post-translational control of HilD. *Mol. Microbiol.* 87, 1045–1060. doi: 10.1111/mmi.12149
- Isberg, R. R., O'Connor, T. J., and Heidtman, M. (2009). The *Legionella pneumophila* replication vacuole: making a cosy niche inside host cells. *Nat. Rev. Microbiol.* 7, 13–24. doi: 10.1038/nrmicro1967
- Jackowski, S., and Rock, C. O. (1983). Ratio of active to inactive forms of acyl carrier protein in *Escherichia coli*. *J. Biol. Chem.* 258, 15186–15191.
- Jackowski, S., and Rock, C. O. (1987). Altered molecular form of acyl carrier protein associated with beta-ketoacyl-acyl carrier protein synthase II (*fabF*) mutants. *J. Bacteriol.* 169, 1469–1473.
- James, B. W., Mauchline, W. S., Dennis, P. J., and Keevil, C. W. (1997). A study of iron acquisition mechanisms of *Legionella pneumophila* grown in chemostat culture. *Curr. Microbiol.* 34, 238–243. doi: 10.1007/s002849900176
- James, B. W., Mauchline, W. S., Dennis, P. J., Keevil, C. W., and Wait, R. (1999). Poly-3-hydroxybutyrate in *Legionella pneumophila*, an energy source for survival in low-nutrient environments. *Appl. Environ. Microbiol.* 2, 822–827.
- James, B. W., Mauchline, W. S., Fitzgeorge, R. B., Dennis, P. J., and Keevil, C. W. (1995). Influence of iron-limited continuous culture on physiology and virulence of *Legionella pneumophila*. *Infect. Immun.* 63, 4224–4230.
- Johnson, W., Varner, L., and Poch, M. (1991). Acquisition of iron by *Legionella pneumophila*: role of iron reductase. *Infect. Immun.* 59, 2376–2381.
- Joshi, A. D., Sturgill-Koszycki, S., and Swanson, M. S. (2001). Evidence that Dot-dependent and -independent factors isolate the *Legionella pneumophila* phagosome from the endocytic network in mouse macrophages. *Cell. Microbiol.* 3, 99–114. doi: 10.1046/j.1462-5822.2001.00093.x
- Keseler, I. M., Collado-Vides, J., Santos-Zavaleta, A., Peralta-Gil, M., Gama-Castro, S., Muniz-Rascado, L., et al. (2011). EcoCyc: a comprehensive database of *Escherichia coli* biology. *Nucleic Acids Res.* 39, D583–D590. doi: 10.1093/nar/gkq1143
- Komander, D., and Rape, M. (2012). The ubiquitin code. *Annu. Rev. Biochem.* 81, 203–229. doi: 10.1146/annurev-biochem-060310-170328
- Kuhle, K., and Flieger, A. (2013). *Legionella* phospholipases implicated in virulence. *Curr. Top. Microbiol. Immunol.* 376, 175–209. doi: 10.1007/82\_2013\_348
- LaRossa, R. A., Van Dyk, T. K., and Smulski, D. R. (1987). Toxic accumulation of alpha-ketobutyrate caused by inhibition of the branched-chain amino acid biosynthetic enzyme acetolactate synthase in *Salmonella typhimurium*. *J. Bacteriol.* 169, 1372–1378.
- Lasbury, M. E., Merali, S., Durant, P. J., Tschang, D., Ray, C. A., and Lee, C. H. (2007). Polyamine-mediated apoptosis of alveolar macrophages during *Pneumocystis pneumonia*. *J. Biol. Chem.* 282, 11009–11020. doi: 10.1074/jbc.M611686200
- Lawhon, S. D., Maurer, R., Suyemoto, M., and Altier, C. (2002). Intestinal short-chain fatty acids alter *Salmonella typhimurium* invasion gene expression and virulence through BarA/SirA. *Mol. Microbiol.* 46, 1451–1464. doi: 10.1046/j.1365-2958.2002.03268.x
- Liles, M. R., Scheel, T. A., and Cianciotto, N. P. (2000). Discovery of a nonclassical siderophore, legiobactin, produced by strains of *Legionella pneumophila*. *J. Bacteriol.* 182, 749–757. doi: 10.1128/JB.182.3.749-757.2000
- Lim, D. B., Oppenheim, J. D., Eckhardt, T., and Maas, W. K. (1987). Nucleotide sequence of the argR gene of *Escherichia coli* K-12 and isolation of its product, the arginine repressor. *Proc. Natl. Acad. Sci. U.S.A.* 84, 6697–6701. doi: 10.1073/pnas.84.19.6697
- Lombard, J., Lopez-Garcia, P., and Moreira, D. (2012). An ACP-independent fatty acid synthesis pathway in archaea: implications for the origin of phospholipids. *Mol. Biol. Evol.* 29, 3261–3265. doi: 10.1093/molbev/mss160
- Lomma, M., Dervins-Ravault, D., Rolando, M., Nora, T., Newton, H. J., Sansom, F. M., et al. (2010). The *Legionella pneumophila* F-box protein Lpp2082 (AnkB) modulates ubiquitination of the host protein parvin B and promotes intracellular replication. *Cell. Microbiol.* 12, 1272–1291. doi: 10.1111/j.1462-5822.2010.01467.x
- Lynch, D., Fieser, N., Glogler, K., Forsbach-Birk, V., and Marre, R. (2003). The response regulator LetA regulates the stationary-phase stress response in *Legionella pneumophila* and is required for efficient infection of *Acanthamoeba castellanii*. *FEMS Microbiol. Lett.* 219, 241–248. doi: 10.1016/S0378-1097(03)00050-8
- Ma, B., Pan, S. J., Zupancic, M. L., and Cormack, B. P. (2007). Assimilation of NAD(+) precursors in *Candida glabrata*. *Mol. Microbiol.* 66, 14–25. doi: 10.1111/j.1365-2958.2007.05886.x
- Maas, W. K. (1994). The arginine repressor of *Escherichia coli*. *Microbiol. Rev.* 58, 631–640.
- Man, W. J., Li, Y., O'Connor, C. D., and Wilton, D. C. (1995). The binding of propionyl-CoA and carboxymethyl-CoA to *Escherichia coli* citrate synthase. *Biochim. Biophys. Acta* 1250, 69–75. doi: 10.1016/0167-4838(95)00044-U
- Maruyama, K., and Kitamura, H. (1985). Mechanisms of growth inhibition by propionate and restoration of the growth by sodium bicarbonate or acetate in *Rhodospseudomonas sphaeroides* S. *J. Biochem.* 98, 819–824.
- McPheat, W. L., Wardlaw, A. C., and Novotny, P. (1983). Modulation of *Bordetella pertussis* by nicotinic acid. *Infect. Immun.* 41, 516–522.
- Melton, A. R., and Weiss, A. A. (1989). Environmental regulation of expression of virulence determinants in *Bordetella pertussis*. *J. Bacteriol.* 171, 6206–6212.
- Mengaud, J. M., and Horwitz, M. A. (1993). The major iron-containing protein of *Legionella pneumophila* is an aconitase homologous with the human iron-responsive element-binding protein. *J. Bacteriol.* 175, 5666–5676.
- Molofsky, A. B., Shetron-Rama, L. M., and Swanson, M. S. (2005). Components of the *Legionella pneumophila* flagellar regulon contribute to multiple virulence traits, including lysosome avoidance and macrophage death. *Infect. Immun.* 73, 5720–5734. doi: 10.1128/IAI.73.9.5720-5734.2005
- Molofsky, A. B., and Swanson, M. S. (2003). *Legionella pneumophila* CsrA is a pivotal repressor of transmission traits and activator of replication. *Mol. Microbiol.* 50, 445–461. doi: 10.1046/j.1365-2958.2003.03706.x
- Molofsky, A. B., and Swanson, M. S. (2004). Differentiate to thrive: lessons from the *Legionella pneumophila* life cycle. *Mol. Microbiol.* 53, 29–40. doi: 10.1111/j.1365-2958.2004.04129.x
- Nasrallah, G. K., Riveroll, A. L., Chong, A., Murray, L. E., Lewis, P. J., and Garduno, R. A. (2011). *Legionella pneumophila* requires polyamines for optimal intracellular growth. *J. Bacteriol.* 193, 4346–4360. doi: 10.1128/JB.01506-10

- Naylor, J., and Cianciotto, N. P. (2004). Cytochrome c maturation proteins are critical for *in vivo* growth of *Legionella pneumophila*. *FEMS Microbiol. Lett.* 241, 249–256. doi: 10.1016/j.femsle.2004.10.028
- Pegg, A. E. (1986). Recent advances in the biochemistry of polyamines in eukaryotes. *Biochem. J.* 234, 249–262.
- Pine, L., George, J. R., Reeves, M. W., and Harrell, W. K. (1979). Development of a chemically defined liquid medium for growth of *Legionella pneumophila*. *J. Clin. Microbiol.* 9, 615–626.
- Poch, M. T., and Johnson, W. (1993). Ferric reductases of *Legionella pneumophila*. *Biomaterials* 6, 107–114. doi: 10.1007/BF00140111
- Pope, C. D., O'Connell, W., and Cianciotto, N. P. (1996). *Legionella pneumophila* mutants that are defective for iron acquisition and assimilation and intracellular infection. *Infect. Immun.* 64, 629–636.
- Price, C. T., Al-Khodori, S., Al-Quadan, T., and Abu Kwaik, Y. (2010). Indispensable role for the eukaryotic-like ankyrin domains of the ankyrin B effector of *Legionella pneumophila* within macrophages and amoebae. *Infect. Immun.* 78, 2079–2088. doi: 10.1128/IAI.01450-09
- Price, C. T., Al-Khodori, S., Al-Quadan, T., Santic, M., Habyarimana, F., Kalia, A., et al. (2009). Molecular mimicry by an F-box effector of *Legionella pneumophila* hijacks a conserved polyubiquitination machinery within macrophages and protozoa. *PLoS Pathog.* 5:e1000704. doi: 10.1371/journal.ppat.1000704
- Price, C. T., Al-Quadan, T., Santic, M., Rosenshine, I., and Abu Kwaik, Y. (2011). Host proteasomal degradation generates amino acids essential for intracellular bacterial growth. *Science* 334, 1553–1557. doi: 10.1126/science.1212868
- Rasis, M., and Segal, G. (2009). The LetA-RsmYZ-CsrA regulatory cascade, together with RpoS and PmrA, post-transcriptionally regulates stationary phase activation of *Legionella pneumophila* Icm/Dot effectors. *Mol. Microbiol.* 72, 995–1010. doi: 10.1111/j.1365-2958.2009.06705.x
- Reeves, M. W., Pine, L., Hutner, S. H., George, J. R., and Harrell, W. K. (1981). Metal requirements of *Legionella pneumophila*. *J. Clin. Microbiol.* 13, 688–695.
- Ristroph, J. D., Hedlund, K. W., and Gowda, S. (1981). Chemically defined medium for *Legionella pneumophila* growth. *J. Clin. Microbiol.* 13, 115–119.
- Robey, M., and Cianciotto, N. P. (2002). *Legionella pneumophila* feoAB promotes ferrous iron uptake and intracellular infection. *Infect. Immun.* 70, 5659–5669. doi: 10.1128/IAI.70.10.5659-5669.2002
- Rolando, M., and Buchrieser, C. (2012). Post-translational modifications of host proteins by *Legionella pneumophila*: a sophisticated survival strategy. *Future Microbiol.* 7, 369–381. doi: 10.2217/fmb.12.9
- Rowbotham, T. J. (1983). Isolation of *Legionella pneumophila* from clinical specimens via amoebae, and the interaction of those and other isolates with amoebae. *J. Clin. Pathol.* 36, 978–986. doi: 10.1136/jcp.36.9.978
- Rowbotham, T. J. (1986). Current views on the relationships between amoebae, legionellae and man. *Isr. J. Med. Sci.* 22, 678–689.
- Sahr, T., Bruggemann, H., Jules, M., Lomma, M., Albert-Weissenberger, C., Cazalet, C., et al. (2009). Two small ncRNAs jointly govern virulence and transmission in *Legionella pneumophila*. *Mol. Microbiol.* 72, 741–762. doi: 10.1111/j.1365-2958.2009.06677.x
- Sanwall, B. D. (1970). Allosteric controls of amphibolic pathways in bacteria. *Bacteriol. Rev.* 34, 20–39.
- Sauer, J. D. (2006). *Contribution of Pht Transporters to Nutrient Acquisition and Differentiation of Intracellular Legionella pneumophila*. Ph.D. thesis, University of Michigan.
- Sauer, J. D., Bachman, M. A., and Swanson, M. S. (2005). The phagosomal transporter A couples threonine acquisition to differentiation and replication of *Legionella pneumophila* in macrophages. *Proc. Natl. Acad. Sci. U.S.A.* 102, 9924–9929. doi: 10.1073/pnas.0502767102
- Schneider, D. R., and Parker, C. D. (1982). Effect of pyridines on phenotypic properties of *Bordetella pertussis*. *Infect. Immun.* 38, 548–553.
- Schnell, J. D., and Hicke, L. (2003). Non-traditional functions of ubiquitin and ubiquitin-binding proteins. *J. Biol. Chem.* 278, 35857–35860. doi: 10.1074/jbc.R300018200
- Serganov, A., and Patel, D. J. (2012). Metabolite recognition principles and molecular mechanisms underlying riboswitch function. *Annu. Rev. Biophys.* 41, 343–370. doi: 10.1146/annurev-biophys-101211-113224
- Sexton, J. A., and Vogel, J. P. (2004). Regulation of hypercompetence in *Legionella pneumophila*. *J. Bacteriol.* 186, 3814–3825. doi: 10.1128/JB.186.12.3814-3825.2004
- Shah, P., and Swiatlo, E. (2008). A multifaceted role for polyamines in bacterial pathogens. *Mol. Microbiol.* 68, 4–16. doi: 10.1111/j.1365-2958.2008.06126.x
- Skaar, J. R., Pagan, J. K., and Pagano, M. (2013). Mechanisms and function of substrate recruitment by F-box proteins. *Nat. Rev. Mol. Cell Biol.* 14, 369–381. doi: 10.1038/nrm3582
- Starai, V. J., and Escalante-Semerena, J. C. (2004). Acetyl-coenzyme A synthetase (AMP forming). *Cell. Mol. Life Sci.* 61, 2020–2030. doi: 10.1007/s00018-004-3448-x
- Starai, V. J., Takahashi, H., Boeke, J. D., and Escalante-Semerena, J. C. (2003). Short-chain fatty acid activation by acyl-coenzyme A synthetases requires SIR2 protein function in *Salmonella enterica* and *Saccharomyces cerevisiae*. *Genetics* 163, 545–555.
- Steinert, M., Emody, L., Amann, R., and Hacker, J. (1997). Resuscitation of viable but nonculturable *Legionella pneumophila* Philadelphia JR32 by *Acanthamoeba castellanii*. *Appl. Environ. Microbiol.* 63, 2047–2053.
- Sturgill-Koszycki, S., and Swanson, M. S. (2000). *Legionella pneumophila* replication vacuoles mature into acidic, endocytic organelles. *J. Exp. Med.* 192, 1261–1272. doi: 10.1084/jem.192.9.1261
- Swanson, M. S., and Isberg, R. R. (1995). Association of *Legionella pneumophila* with the macrophage endoplasmic reticulum. *Infect. Immun.* 63, 3609–3620.
- Swanson, M. S., and Isberg, R. R. (1996). Identification of *Legionella pneumophila* mutants that have aberrant intracellular fates. *Infect. Immun.* 64, 2585–2594.
- Sze, C. W., and Li, C. (2011). Inactivation of bb0184, which encodes carbon storage regulator A, represses the infectivity of *Borrelia burgdorferi*. *Infect. Immun.* 79, 1270–1279. doi: 10.1128/IAI.00871-10
- Tabor, C. W., and Tabor, H. (1985). Polyamines in microorganisms. *Microbiol. Rev.* 49, 81–99.
- Tabor, H., Hafner, E. W., and Tabor, C. W. (1980). Construction of an *Escherichia coli* strain unable to synthesize putrescine, spermidine, or cadaverine: characterization of two genes controlling lysine decarboxylase. *J. Bacteriol.* 144, 952–956.
- Tesh, M. J., and Miller, R. D. (1981). Amino acid requirements for *Legionella pneumophila* growth. *J. Clin. Microbiol.* 13, 865–869.
- Tesh, M. J., Morse, S. A., and Miller, R. D. (1983). Intermediary metabolism in *Legionella pneumophila*: utilization of amino acids and other compounds as energy sources. *J. Bacteriol.* 154, 1104–1109.
- Traxler, M. F., Summers, S. M., Nguyen, H. T., Zacharia, V. M., Hightower, G. A., Smith, J. T., et al. (2008). The global, ppGpp-mediated stringent response to amino acid starvation in *Escherichia coli*. *Mol. Microbiol.* 68, 1128–1148. doi: 10.1111/j.1365-2958.2008.06229.x
- VanRheenen, S. M., Luo, Z. Q., O'Connor, T., and Isberg, R. R. (2006). Members of a *Legionella pneumophila* family of proteins with ExoU (phospholipase A) active sites are translocated to target cells. *Infect. Immun.* 74, 3597–3606. doi: 10.1128/IAI.02060-05
- Vergara-Irigaray, N., Chavarri-Martinez, A., Rodriguez-Cuesta, J., Miller, J. F., Cotter, P. A., and Martinez de Tejada, G. (2005). Evaluation of the role of the Bvg intermediate phase in *Bordetella pertussis* during experimental respiratory infection. *Infect. Immun.* 73, 748–760. doi: 10.1128/IAI.73.2.748-760.2005
- Viner, R., Chetrit, D., Ehrlich, M., and Segal, G. (2012). Identification of two *Legionella pneumophila* effectors that manipulate host phospholipids biosynthesis. *PLoS Pathog.* 8:e1002988. doi: 10.1371/journal.ppat.1002988
- Viswanathan, V. K., Edelstein, P. H., Pope, C. D., and Cianciotto, N. P. (2000). The *Legionella pneumophila* iraAB locus is required for iron assimilation, intracellular infection, and virulence. *Infect. Immun.* 68, 1069–1079. doi: 10.1128/IAI.68.3.1069-1079.2000
- Viswanathan, V. K., Kurtz, S., Pedersen, L. L., Abu-Kwaik, Y., Krcmarik, K., Mody, S., et al. (2002). The cytochrome c maturation locus of *Legionella pneumophila* promotes iron assimilation and intracellular infection and contains a strain-specific insertion sequence element. *Infect. Immun.* 70, 1842–1852. doi: 10.1128/IAI.70.4.1842-1852.2002
- Wieland, H., Ullrich, S., Lang, F., and Neumeister, B. (2005). Intracellular multiplication of *Legionella pneumophila* depends on host cell amino acid transporter SLC1A5. *Mol. Microbiol.* 55, 1528–1537. doi: 10.1111/j.1365-2958.2005.04490.x
- Williams, C. L., and Cotter, P. A. (2007). Autoregulation is essential for precise temporal and steady-state regulation by the *Bordetella* BvgAS phosphorelay. *J. Bacteriol.* 189, 1974–1982. doi: 10.1128/JB.01684-06
- Wolfe, A. J. (2005). The acetate switch. *Microbiol. Mol. Biol. Rev.* 69, 12–50. doi: 10.1128/MMBR.69.1.12-50.2005
- Wolfe, A. J. (2010). Physiologically relevant small phosphodonors link metabolism to signal transduction. *Curr. Opin. Microbiol.* 13, 204–209. doi: 10.1016/j.mib.2010.01.002

- Wolfe, A. J., Chang, D. E., Walker, J. D., Seitz-Partridge, J. E., Vidaurri, M. D., Lange, C. F., et al. (2003). Evidence that acetyl phosphate functions as a global signal during biofilm development. *Mol. Microbiol.* 48, 977–988. doi: 10.1046/j.1365-2958.2003.03457.x
- Xu, L., Shen, X., Bryan, A., Banga, S., Swanson, M. S., and Luo, Z. Q. (2010). Inhibition of host vacuolar H<sup>+</sup>-ATPase activity by a *Legionella pneumophila* effector. *PLoS Pathog.* 6:e1000822. doi: 10.1371/journal.ppat.1000822
- Xu, X., Liu, Z., Fang, M., Yu, H., Liang, X., Li, X., et al. (2012). *Helicobacter pylori* CagA induces ornithine decarboxylase upregulation via Src/MEK/ERK/c-Myc pathway: implication for progression of gastric diseases. *Exp. Biol. Med.* 237, 435–441. doi: 10.1258/ebm.2011.011199
- Yin, Y., He, X., Szwedczyk, P., Nguyen, T., and Chang, G. (2006). Structure of the multidrug transporter EmrD from *Escherichia coli*. *Science* 312, 741–744. doi: 10.1126/science.1125629
- Yip, E. S., Burnside, D. M., and Cianciotto, N. P. (2011). Cytochrome c4 is required for siderophore expression by *Legionella pneumophila*, whereas cytochromes c1 and c5 promote intracellular infection. *Microbiology* 157, 868–878. doi: 10.1099/mic.0.046490-0
- Zhao, B., and Houry, W. A. (2010). Acid stress response in enteropathogenic gammaproteobacteria: an aptitude for survival. *Biochem. Cell Biol.* 88, 301–314. doi: 10.1139/O09-182
- Zheng, H., Chatfield, C. H., Liles, M. R., and Cianciotto, N. P. (2013). Secreted pyomelanin of *Legionella pneumophila* promotes bacterial iron uptake and growth under iron-limiting conditions. *Infect. Immun.* 81, 4182–4191. doi: 10.1128/IAI.00858-13
- Conflict of Interest Statement:** The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.
- Received: 31 December 2013; accepted: 23 January 2014; published online: 11 February 2014.
- Citation: Fonseca MV and Swanson MS (2014) Nutrient salvaging and metabolism by the intracellular pathogen *Legionella pneumophila*. *Front. Cell. Infect. Microbiol.* 4:12. doi: 10.3389/fcimb.2014.00012
- This article was submitted to the journal *Frontiers in Cellular and Infection Microbiology*.
- Copyright © 2014 Fonseca and Swanson. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) or licensor are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.